Mapping

胃

担当 小坂 太郎

白色光
(4枚)

TXI

20

1234

インジゴカルミン散布

NBI
(7枚)

ESD

マーキング

ESD後

口側

肛門側

口側

肛門側

口側

肛門側

口側

肛門側

口側

肛門側

口側

肛門側

12

1

口側

肛門側

口側

肛門側

口側

肛門側

12

11

10

9

8

7

6

5

4

3

2

1

口側

肛門側

口側

肛門側

口側

肛門側

口側

肛門側

key slice

口側

12

11

10

9

8

7

6

5

4

3

2

1

肛門側

口側

肛門側

口側

肛門側

12

11

10

9

8

7

6

5

4

3

2

1

口側

12

11

10

9

8

7

6

5

4

3

2

1

肛門側

口側

肛門側

12

11

10

9

8

7

6

5

4

3

2

1

口側

肛門側

12

11

10

9

8

7

6

5

4

3

1

2

口側

肛門側

12

11

10

9

8

7

6

5

4

3

2

1

口側

12

11

10

9

8

7

6

5

4

3

2

1

肛門側

口側

肛門側

12

11

10

9

8

7

6

5

2

1

口側

12

11

10

9

8

7

6

5

4

3

2

1

肛門側

口側

12

11

10

9

8

7

6

5

4

3

2

1

肛門側

口側

肛門側

12

11

10

9

8

7

6

5

4

3

2

1

口側

12

11

10

9

8

7

6

5

4

3

2

1

肛門側

口側

12

11

10

9

8

7

6

5

4

3

2

1

肛門側

口側

肛門側

12

11

10

9

8

7

6

5

4

3

2

1

口側

12

11

10

9

8

7

6

5

4

3

2

1

肛門側

口側

肛門側

12

11

10

9

8

7

6

5

4

3

2

1

病理診断

tubular adenocarcinoma,
moderately differentiated(tub2>tub1>muc),
T1b1, SM1, early, UL1, Ly0, V0, HM0, VM0.
eCure B

mapping

口側

肛門側

口側

肛門側

口側

肛門側

12

11

10

9

8

7

6

5

4

3

2

1

口側

肛門側

12

11

10

9

8

7

6

5

4

3

2

1

— M

— SM1

口側

— M

— SM1

肛門側

口側

肛門側

12
11
10
9
8
7
6
5
4
3
2
1

tub1
tub2
muc

口側

肛門側

- tub1
- tub2
- muc

口側

肛門側

— pUL1

12

11

10

9

8

7

6

5

4

3

2

1

口側

肛門側

— pUL1

口側

肛門側

口側

肛門側

口側

12

1

11

2

10

3

9

4

8

5

7

6

肛門側

口側

肛門側

M

SM1

口側

肛門側

— M

— SM1

口側

12

11

10

9

8

7

6

5

4

3

2

1

tub1

tub2

muc

肛門側

口側

肛門側

- tub1
- tub2
- muc

ESDの適応

壁深達度	潰瘍	分化型		未分化型	
		≤ 2 cm	> 2 cm	≤ 2 cm	> 2 cm
cT1a (M)	UL0	★			
	UL1				
cT1b (SM)

EMR/ESDの絶対適応病変
 ESDの絶対適応病変
相対適応病変

cT1a (M) : 粘膜内癌 (術前診断), cT1b (SM) : 粘膜下層浸潤癌 (術前診断)
 UL : 潰瘍 (癒痕) 所見
 Figure 1 腫瘍因子による適応の分類.

胃癌に対するESD/EMRガイドライン (第2版)

- ① UL0, cT1a, 分化型癌
- ② UL0, cT1a, 未分化癌
- ③ 3cm以下, UL1, cT1a, 分化型癌

ESDの適応

壁深達度	潰瘍	分化型		未分化型	
		≤ 2 cm	> 2 cm	≤ 2 cm	> 2 cm
cT1a (M)	UL0	★			
	UL1	■			
cT1b (SM)					

cT1a (M) : 粘膜内癌 (術前診断), cT1b (SM) : 粘膜下層浸潤癌 (術前診断)
 UL : 潰瘍 (癒痕) 所見
 Figure 1 腫瘍因子による適応の分類.

胃癌に対するESD/EMRガイドライン (第2版)

- ① UL0, cT1a, 分化型癌
- ② UL0, cT1a, 未分化癌
- ③ 3cm以下, UL1, cT1a, 分化型癌

内視鏡的根治度(eCura)

壁深達度	潰瘍	分化型	未分化型
pT1a (M)	UL0		≤ 2 cm > 2 cm
	UL1	≤ 3 cm > 3 cm	
pT1b1 (SM1)		≤ 3 cm > 3 cm	
pT1b2 (SM2)

eCureA*

eCureB*

eCureC-2

胃癌に対するESD/EMRガイドライン (第2版)

eCuraA : 絶対適応病変で、HM0, VM0, Ly0, V0

eCuraB : 3cm以下の分化型癌、SM1

eCureC : それ以外 (非治癒切除)

内視鏡的根治度(eCura)

壁深達度	潰瘍	分化型	未分化型
pT1a (M)	UL0		≤ 2 cm > 2 cm
	UL1	≤ 3 cm > 3 cm
pT1b1 (SM1)		≤ 3 cm > 3 cm
pT1b2 (SM2)

eCuraA*

eCuraB*

eCuraC-2

胃癌に対するESD/EMRガイドライン (第2版)

eCuraA : 絶対適応病変で、HM0,VM0,Ly0,V0

eCuraB : 3cm以下の分化型癌、SM1

eCuraC : それ以外 (非治癒切除)

eCuraBについて

Original article

Incidence of lymph node metastasis from early gastric cancer: estimation with a large number of cases at two large centers

TAKUJI GOTODA¹, AKIO YANAGISAWA², MITSURU SASAKO³, HIROYUKI ONO¹, YUKIHIRO NAKANISHI⁴,
TADAKAZU SHIMODA⁵, and YO KATO²

¹Department of Endoscopy, National Cancer Center Hospital, Tokyo, Japan

²Department of Pathology, Cancer Institute, Tokyo, Japan

³Department of Surgical Oncology, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

⁴Pathology Division, National Cancer Center Research Institute, Tokyo, Japan

⁵Department of Clinical Laboratory, National Cancer Center Hospital, Tokyo, Japan

- 胃切除術を施行された5265例
 - M癌：3016例（リンパ節転移65例）
 - SM癌：2249例（リンパ節転移402例）

に対して、リンパ節転移（LNM）のリスク因子を検討

Table 3. Relationship between clinicopathological factors and lymph node metastasis in submucosal invasive cancer; univariate analysis results

	Total	Status of LN metastasis			<i>P</i> value
		Negative	Positive	Percent	
Sex					
M	1424	1173	251	17.6	0.0215
F	629	491	138	21.9	
Tumor location					
U	619	501	118	19.1	0.9402
M	1028	836	192	18.7	
L	406	327	79	19.5	
Macroscopic type					
Elevated	440	331	109	24.8	0.0003
Depressed	1505	1247	258	17.1	
Tumor size					
≤ 10 mm	114	105	9	7.9	<0.0001
≤ 20 mm	512	444	68	13.3	
≤ 30 mm	677	572	105	15.5	
> 31 mm	946	726	220	23.3	
Histological type					
Differentiated	1277	1060	217	17.0	0.2110
Undifferentiated	972	787	185	19.0	
Ulcer findings					
Absence	790	640	150	19.0	0.9712
Presence	1263	1024	239	18.9	
Lymphatic-vascular involvement					
Absence	1350	1218	132	9.8	<0.0001
Presence	703	446	257	36.6	
Degree of submucosal penetration					
SM1	364	333	32	8.8	<0.0001
SM2	1272	971	301	23.7	

SM1, Penetration of submucosal layer less than 500 μ m from muscularis mucosa; SM2, penetration of 500 μ m or more

Table 3. Relationship between clinicopathological factors and lymph node metastasis in submucosal invasive cancer; univariate analysis results

	Total	Status of LN metastasis			P value
		Negative	Positive	Percent	
Sex					
M	1424	1173	251	17.6	0.0215
F	629	491			
Tumor location					
U	619	501			<div style="border: 2px solid red; padding: 5px;"> <ul style="list-style-type: none"> ▪ 腫瘍径 > 31mm ▪ Ly1, V1 ▪ SM2 <p>で有意にリンパ節転移率上昇</p> </div>
M	1028	836			
L	406	327			
Macroscopic type					
Elevated	440	331			0.2110
Depressed	1505	1247			
Tumor size					
≤10 mm	114	105			0.9712
≤20 mm	512	444	68	13.3	
≤30 mm	677	572	105	15.5	
>31 mm	946	726	220	23.3	
Histological type					
Differentiated	1277	1060	217	17.0	0.2110
Undifferentiated	972	787	185	19.0	
Ulcer findings					
Absence	790	640	150	19.0	0.9712
Presence	1263	1024	239	18.9	
Lymphatic-vascular involvement					
Absence	1350	1218	132	9.8	<0.0001
Presence	703	446	257	36.6	
Degree of submucosal penetration					
SM1	364	333	32	8.8	<0.0001
SM2	1272	971	301	23.7	

SM1, Penetration of submucosal layer less than 500μm from muscularis mucosa; SM2, penetration of 500μm or more

Table 3. Relationship between clinicopathological factors and lymph node metastasis in submucosal invasive cancer; univariate analysis results

	Total	Status of LN metastasis			P value
		Negative	Positive	Percent	
Sex					
M	1424	1173	251	17.6	0.0215
F	629	491	138	21.9	
Tumor location					
U	619	501	118	19.1	0.9402
M	1028	836	192	18.7	
L	406	327	79	19.5	
Macroscopic type					
Elevated	440	331	109	24.8	0.0003
Depressed	1505	1247	258	17.1	
Tumor size					
≤10 mm	114	105	9	7.9	<0.0001
≤20 mm	512	444	68	13.3	
≤30 mm	677	572	105	15.5	
>31 mm	946	726	220	23.3	
Histological type					
Differentiated	1277	1060	217	17.0	
Undifferentiated	972	787	185	19.0	
Ulcer findings					
Absence	790	640	150	19.0	0.9712
Presence	1263	1024	239	18.9	
Lymphatic-vascular involvement					
Absence	1350	1218	132	9.8	<0.0001
Presence	703	446	257	36.6	
Degree of submucosal penetration					
SM1	364	333	32	8.8	<0.0001
SM2	1272	971	301	23.7	

ULの有無では
リンパ節転移に有意差なし

SM1, Penetration of submucosal layer less than 500μm from muscularis mucosa; SM2, penetration of 500μm or more

- SM浸潤癌で、リンパ節転移の確率が高いもの
 - 腫瘍径 > 30mm
 - Ly1
 - V1
 - SM2
- ULの有無ではリンパ節転移に有意差なし

Lymph node metastasis by tumor size in SM1 differentiated tumors without lymphatic-vascular involvement

	Number	LNМ	%
≦10mm	28	0	0.0
≦20mm	59	0	0.0
≦30mm	58	0	0.0
>31mm	78	2	2.6
Total	223	2	0.9

- SM1, ly0, v0で腫瘍径≦30mmであれば、リンパ節転移は0%

eCuraB症例における

5年全生存率 87.4%, 10年疾患特異的生存率 96.8%

胃癌再発2例、異時多発による死亡2例

STOMACH

Long-term Survival After Endoscopic Resection For Gastric Cancer: Real-world Evidence From a Multicenter Prospective Cohort

Haruhisa Suzuki,¹ Hiroyuki Ono,² Toshiaki Hirasawa,³ Yoji Takeuchi,⁴ Kenji Ishido,⁵ Shu Hoteya,⁶ Tomonori Yano,⁷ Shinji Tanaka,⁸ Yosuke Toya,⁹ Masahiro Nakagawa,¹⁰ Takashi Toyonaga,¹¹ Kenichi Takemura,¹² Kingo Hirasawa,¹³ Mitsuru Matsuda,¹⁴ Hironori Yamamoto,¹⁵ Yosuke Tsuji,¹⁶ Satoru Hashimoto,¹⁷ Maeda Yuki,² Tsuneo Oyama,¹⁸ Ryuta Takenaka,¹⁹ Yoshinobu Yamamoto,²⁰ Yuji Naito,²¹ Katsumi Yamamoto,²² Nozomu Kobayashi,²³ Yoshiro Kawahara,²⁴ Masaaki Hirano,²⁵ Shigeto Koizumi,²⁶ Shinichiro Hori,²⁷ Masahiro Tajika,²⁸ Takuto Hikichi,²⁹ Kenshi Yao,³⁰ Chizu Yokoi,³¹ Ken Ohnita,³² Yasuhiro Hisanaga,³³ Tetsuya Sumiyoshi,³⁴ Shinji Kitamura,³⁵ Hisao Tanaka,³⁶ Ryo Shimoda,³⁷ Taichi Shimazu,³⁸ Kohei Takizawa,² Satoshi Tanabe,³⁹ Hitoshi Kondo,³⁴ Hiroyasu Iishi,^{4,40} Motoki Ninomiya,⁴¹ and Ichiro Oda,¹ on behalf of the J-WEB/EGC group

J-WEB/EGC

- 早期胃癌に対する内視鏡治療9054症例
- 全国多施設での前向きコホート研究
- 5年全生存率, 5年疾患特異的生存率を調査

Long-term Survival after Endoscopic Resection for Gastric Cancer: Real-world Evidence from a Multicenter Prospective Cohort

9054 Early Gastric Cancer Patients
from Japanese Multicenter Cohort

5-year Overall Survival
89.0% (95% CI, 88.3%-89.6%)

HRs for all-cause mortality

Curability of Gastric Endoscopic Resection	n	Multivariate-HRs (95% CI)
Category A1: Differentiated-type, pT1a, UL (-), ≤2 cm	4545	REF
Category A2: Differentiated-type, pT1a, UL (-), >2 cm or UL (+), ≤3 cm	2084	1.03 (0.87-1.21)
Category A3: Undifferentiated-type, pT1a, UL (-), ≤2 cm	226	1.18 (0.68-2.07)
Category B: Differentiated-type, pT1b (SM1), ≤3 cm	387	1.09 (0.80-1.49)
Category C: Noncurative resection	1812	1.41 (1.21-1.65)

0.5 1 2

Clinical Gastroenterology and Hepatology

Endoscopic Resection

- 内視鏡的根治度ごとの5年生存率を比較

壁深達度	潰瘍	分化型		未分化型	
		≤ 3 cm	> 3 cm	≤ 2 cm	> 2 cm
pT1a (M)	UL0	A1	A2	A3	
	UL1	A2			
pT1b1 (SM1)		B			
pT1b2 (SM2)

eCureA*

eCureB*

eCureC-2

Table 3. HRs for All-cause Mortality

	n	Death	Person year	Age and sex-adjusted HRs		
				Univariate HRs HR (95% CI)	HRs HR (95% CI)	Multivariate-adjusted HRs HR (95% CI)
Curability						
Category A1	4545	433	21292	1	1	1
Category A2	2084	213	9587	1.09 (0.93-1.29)	1.03 (0.87-1.21)	1.03 (0.87-1.21)
Category A3	226	13	1076	0.59 (0.34-1.02)	1.44 (0.83-2.52)	1.18 (0.68-2.07)
Category B	387	44	1794	1.21 (0.88-1.64)	1.07 (0.79-1.46)	1.09 (0.80-1.49)
Category C	1812	263	8258	1.57 (1.35-1.83)	1.43 (1.23-1.67)	1.41 (1.21-1.65)

• 全死因死亡率のHRは、eCureA,Bで有意差なし

まとめ

- eCuraBにおいても、eCuraAと同等のリンパ節転移率、長期予後が期待できる。